REFERENCES Crozier, W. J., 1924-25, J. Gen. Physiol., 7, 189. Mast, S. O., 1914, Bull. Bur. Fish., 34, 173. Parker, G. H., 1936, "Color Changes of Animals in Relation to Nervous Activity." (Chap. 3.) Philadelphia, Pa. Sumner, F. B., 1911, J. Exp. Zool., 10, 409.

# PRELIMINARY REPORT ON A NEW TUBULARIAN HYDROID

SAMUEL STOCKTON MILES Princeton University\*

A single species of an apparently hitherto undescribed genus of Tubularian hydroid was found in the waters off the Mount Desert



Island Biological Laboratory. (See Fig. 1.) It is a solitary mud-living form taken at between forty and sixty feet. The hydranth has a whorl of 6 to 8 oral tentacles and a whorl of 6 to 16 basal tentacles, the latter with rings of nematocysts along their length. Below the hydrauth the foot or tail is enveloped in a perisarc which is continued posteriorly in an irregular sticky sheath containing small ovoid bodies. The length of the hydroid exclusive of this sheath is about 5 mm. The medusae are formed in the region between the whorls of tentacles and have one tentacle in the immature stage observed.

FIG. 1.—The hydroid in expanded condition.

\* Now of The Johns Hopkins University Medical School.

The ovoid bodies in the posterior portion of the perisarcal sheath are formed by successive constrictions of the foot. These develop into young hydroids directly or after one or more transverse fissions.

Because of the arrangement of the tentacles, the position of development of the medusae, the solitary nature of this hydroid and the single tentacle of the medusae it evidently belongs in the family, *Corymorphidae*. The generic name *Dahlgrenella*, nov. gen., is chosen to express my gratitude to Professor Ulric Dahlgren for his kind assistance and for the method of collecting mud-living hydroids devised by him. The specific name, *farcta*, nov. sp., is proposed as descriptive of the perisarc with its ovoid reproductive bodies, being from the Latin verb "to stuff" from which the Latin "farcimen" for sausage is derived.

#### A STUDY OF THE ACTION OF CERTAIN DRUGS ON THE CIRCULATION OF THE DOGFISH

## J. T. HALSEY and ERNEST M. EVANS Tulane University and University of Pennsylvania

Ergotamine (as Gynergen, Sandoz) neither reversed nor diminished the augmentor effect of adrenalin on the blood-pressure measured in the ventral aorta of the dogish, *Squalus acanthias*.

Previous injections of cocain did not sensitize to adrenalin nor desensitize to ephedrin.

Posterior pituitary extract (as Pituitrin, P. D. & Co.), and oxytocin (as Pitocin, P. D. & Co.) and vasopressin (as Pitressin, P. D. & Co.) all exerted an augmentor effect on the blood-pressure in the ventral aorta. No depressor effect was exerted by subsequent injections. The effects on the heart rate were variable, usually no significant change resulting.

Chlorazol-fast-pink, an azodye, which, in dosage of 80 to 100 mg. per kilo, has been successfully used in rabbits, cats and dogs as an anticoagulant in vivo, exerts, in much smaller dosage (20 to 30 mg. per kilo), so depressing an action on the circulation of the dogfish, that its use as an agent to prevent clotting in the arterial cannulae had to be abandoned.

## TISSUE CULTURE STUDIES ON DOGFISH PITUITARY

## LELAND B. RANSOM Rutgers University\*

Following the work of M. R. and W. H. Lewis in 1935, the hypophysis of *Squalus acanthias* was again grown successfully in tissue culture. Various media were employed, the type of growth was

\* Now of The Johns Hopkins University.