

Using diameter of the test as the measure of age, the condition of the ciliates was studied in 152 urchins (diam. 8-65 mm.) which represented 6 age-groups (1.5-6.5 years, in increments of 1 year). Urchins 8 mm. in diam. (age about 1.5 years) were uninfected; urchins 9-14 mm. (age likewise 1.5 years) were lightly infected (1-28 ciliates of each species per urchin), but dividing individuals were absent. Urchins 15-23 mm. (age about 2.5 years) contained large numbers of dividing specimens of all 3 species. All the urchins of the remaining age-groups were infected, and *B. gracilis* was dividing in every urchin, but the division of *E. borealis* and *M. indomita* was sporadic and could not be correlated with the age of the host.

The results indicate that urchins become infected in their second summer (at age 1.5 years) and that their ciliate populations build up rapidly in the third summer. Once the populations are established, infrequent division suffices to maintain *E. borealis* and *M. indomita*, though constant division is necessary to maintain *B. gracilis*, which, unlike its confreres, inhabits the rectum and escapes regularly among the fecal pellets.

Further Observations on the Relation of the Ciliate (*Conchophthirus mytili*) to Its Host, (*Mytilus edulis*)

C. Dale Beers
University of North Carolina

Various studies on the behavior of metazoan commensals have shown that some commensals respond positively to their respective hosts or even to host fragments (Davenport, J. Mar. Biol. Assoc. U. K. 32:161, 1953). Such studies raise the question as to whether *Conchophthirus mytili*, an ectocommensal of the foot of the mussel *Mytilus edulis*, displays similar responses.

Since *C. mytili* tolerates pure sea water reasonably well (Beers, J. Elisha Mitchell Sci. Soc. 75:3, 1959), specimens were washed from the foot of freshly collected mussels into "outside" sea water, meaning water collected well beyond the mussel beds, or into artificial sea water (M. B. L. formula). Their response was tested in turn to the detached whole foot and to excised pieces of living foot, gill, mantle and labial palp, but the results were negative. However, specimens that arrived by chance on the epithelial surface of the foot or specimens that were placed on it by means of a pipette remained on it for periods that varied from 9 to 12 hours. But specimens that arrived on the other surfaces that have been mentioned usually remained no longer than a minute before swimming away.

The results indicate that *C. mytili* is carried passively through the incurrent siphon into the mantle chamber of the host and that its arrival on the foot is dictated by chance. Once on the foot, however, it remains "bound" to the epithelial surface of the foot, in all probability by a chemical attractant, though the nature of the attractant is undetermined.