

coloration and turned silver, resembling the silver hue of eels spontaneously migrating to the sea. Freshwater eels prepared by 10 days of hydrocortisone injections (400 μ g/100 gm/day) withstood sudden transfer to full strength seawater without preliminary residence in 50% seawater, and their plasma chloride did not rise as high as that of control eels transferred abruptly to full strength seawater (W. MacKay, report #17 this issue).

The data indicate that hydrocortisone induces a series of changes in freshwater eels, including a rise in Na-K-ATPase of gill and intestine, that successfully prepares these euryhaline teleosts to combat the osmotic stress of migration to seawater.

1969 #7

RELATION OF SALT AND WATER TRANSPORT TO FINE STRUCTURE OF THE INTESTINE AND GILLS OF THE EEL, Anguilla rostrata AND THE DOGFISH, Squalus acanthias

W. L. Doyle, The University of Chicago, Chicago, Ill.

(1) The spiral valve of the dogfish was isolated and the lumen and artery perfused with saline to establish conditions of maximal and minimal rates of water transport across the mucosa. Under such conditions the morphological configuration of the intercellular space in the epithelia should show correlative change. An additional factor is the intraluminal hydrostatic pressure and the experimental procedure was designed to determine the effects of changes in this parameter. Specimens of these preparations were fixed and embedded for electron microscopy. They are currently being studied.

(2) The transfer of eels from fresh to salt water is accompanied by increased salt and water transport in the gills and gut. In collaboration with Franklin Epstein, Ralph Janicki, and William MacKay samples were obtained for electron microscopy of the gills and the proximal gut to provide morphological correlates of the functional states observed by them. The existence of elevated Na-KATPase levels in gills of fresh water eels treated with hydrocortisone (see Epstein this bulletin) should provide interesting comparisons with corresponding levels of enzyme activity and the morphological changes usually associated with animals adapted to sea water.

(3) Preliminary observations on the fine structure of mesothelia found on dogfish mesentery indicate the presence of junctional complexes of the kind found in gill epithelia (close junction, zonula adherens and maculae densa). Other aspects of the fine structure may provide clues to the low rates of absorption of drugs and metabolites from the peritoneal cavity in this animal.

This work was supported by a grant from the National Science Foundation, GB 8312.

1969 #8

THE TRANSPORT OF VARIOUS ORGANIC MATERIALS FROM BLOOD TO TISSUE IN Squalus acanthias

Joseph Fenstermacher, Andres Roomet, Jeffrey Ratner, and David P. Rall, National Cancer Institute, Bethesda, Md. 20014

Studies of the blood to tissue transport of eight organic compounds have been undertaken in