

William Mackay samples were obtained for electron microscopy of the gills and proximal gut to provide morphological correlates of the functional states observed by them. (See this Bulletin Volume 9.) The existence of elevated Na-K ATPase levels in gills of fresh water eels treated with cortisol provides opportunity to study cytological changes in chloride cells due to hormonal influences in the absence of changes due to transfer to saline waters. Under experimental conditions the branching tubular reticulum in the chloride cells varies from an open three dimensional network in a pale cytoplasm to a system so closely branched that the cytoplasm surrounding it is reduced to a continuum of similar dimensions. In the latter case the cytoplasm is dense. There is wide variation in the cytology of chloride cells among individual eels, especially in those from fresh water, and the variability is pronounced for eels in intermediate stages of adaptation or hormone treatment. Despite these variations and the difficulties of making adequate surveys at the electron microscopic level there appears to be a general progression in morphological elaboration of the tubular system corresponding to the duration of treatment with cortisol or adaptation to sea water. With cortisol treatment, especially at intermediate stages (3 and 7 days) there are many cells which appear to be differentiating. They show active nucleoli, increased perinuclear granular endoplasmic reticulum and patches of distended, closely branching tubular reticulum resembling fenestrated cisternae. Other reports have indicated an increased number of chloride cells several days after transfer from fresh to salt water and a similar increase may be expected in eels. Chloride cells require special procedures for recognition at the light microscopy level and such methods have not distinguished the differentiating cells reported here.

It would appear that the amount of tubular endoplasmic reticulum in the chloride cells of the eel gill and the number of chloride cells both increase during adaptation to sea water and as a result of treatment of fresh water eels with cortisol.

Our fresh water eels have varied considerably in size (age) and natural hormonal conditions (yellow to silver). We have the impression that the number of swollen and abnormal cells is greatest in fresh water eels taken late in the summer. Under the stimulus of adaptation to sea water or treatment with cortisol these cells are replaced by newly differentiated cells. Eels obtained in 1970 from mixed sources, kept for a while in brackish water and subsequently readapted to fresh water showed fewer swollen and deteriorating cells than those obtained from fresh water eels late in the season.

Supported by a grant (GB 8312) from the National Science Foundation.

1970 #7

OCCURRENCE OF A VIRUS IN ERYTHROCYTES OF THE EEL, Anguilla

William L. Doyle, The University of Chicago, Chicago, Ill.

In electron microscopic sections of the gill and intestine of one specimen of the eel, Anguilla rostrata, it was observed that about twenty percent of the erythrocytes had clusters of virus particles in the cytoplasm of these nucleated, mitochondria-bearing blood cells. In sections the particles are hexagonal in outline with a diameter of about 0.16 microns. The circular core has a diameter of about 0.10 microns and is variable in intensity of staining with uranyl and lead. This occurrence seems sufficiently unusual to merit noting.