

As may be seen from Figure 2, the clearance ratio of methylamine to PEG also declined with time. However, in this case, the clearance ratio was considerably greater than one which indicates that methylamine is actively secreted. The clearance ratio from three fish ranged from thirty to seven.

The mean ratio of urine flow for these fish was 0.23 ± 0.05 ml/hr, based on fourteen clearance periods. The mean clearance of PEG was 0.28 ± 0.07 ml/hr. The tissue to plasma ratio of methylamine to that for PEG, determined for one fish, was 16.49 ± 0.51 ($n = 5$).

The data thus suggests that methylamine is secreted into the lumen by the transport system for weak (organic) cations (see, e.g., Am. J. Physiol. 225:1123, 1973).

This study was supported by NIH grant AM 12619-10 and the Whitehall Foundation. The assistance of Mr. Jonathan Goldstein was appreciated.

THE RENAL HANDLING OF D-FRUCTOSE BY THE WINTER FLOUNDER (*Pseudopleuronectes americanus*)

G. Booz, D. E. Pew and A. Kleinzeller, University of Pennsylvania, Dept. of Physiology, Philadelphia, Pennsylvania

The renal handling of D-fructose by the winter flounder was characterized employing the teased tubule preparation and clearance studies.

The *in vitro* studies, using teased renal tubules, were carried out as previously described (Kleinzeller et al., Am. J. Physiol. 232:F227, 1977). The tissue to medium ratio (T/M) for seven pieces of tissue, incubated for 60 minutes in saline having an initial concentration of 1 mM D-fructose, was 0.61 ± 0.02 . (All values are expressed as mean \pm the standard error.) No significant difference in the T/M was observed over a concentration range of 0.05 to 5.0 mM; i.e., saturation of a carrier site could not be demonstrated. Glucose, 2-deoxyglucose and mannose, which have previously been shown to share the same carrier at the antiluminal face of the cell, did not inhibit the uptake of D-fructose when present at a fivefold molar excess. Also, 0.5 mM phlorizin, phloretin or ouabain did not significantly reduce the T/M value.

Clearance studies were carried out as previously described (Am. J. Physiol. 231:603, 1976). The clearance of D-fructose was not significantly different from the GFR (clearance of PEG) at a plasma fructose range of 50 to 100 μ M. The mean clearance ratio obtained from six fish was 1.05 ± 0.06 , based on 29 clearance periods. No correlation was observed between either the GFR or the rate of urine flow and the clearance of D-fructose. An additional load of D-glucose, estimated to be 1 mM, had no effect on the clearance of D-fructose.

In brief, no active reabsorption of D-fructose by the winter flounder kidney was observed. Judging from the results obtained with teased tubules, fructose enters the cells at the antiluminal face by a pathway which is not carrier mediated. This investigation was supported in part by USPHS grant AM 12619 and the Whitehall Foundation. The assistance of Mr. Jonathan Goldstein was appreciated.

THE HANDLING OF D-MANNOSE AT THE BRUSH BORDER OF THE KIDNEY IN THE WINTER FLOUNDER (*Pseudopleuronectes americanus*)

G. Booz,* J. Goldstein, J. Pritchard** and A. Kleinzeller,* *Dept. of Physiology, University of Pennsylvania, Philadelphia, Pennsylvania; **National Institute of Environmental Health, NIH, Research Triangle Park, North Carolina

Studies on the handling of D-mannose by the winter flounder (*Pseudopleuronectes americanus*) were continued in view of persistent reports (Silverman, Biochim. Biophys. Acta 457:303, 1976) that in the brush border of the mammalian kidney a specific carrier is responsible for the phlorizin-insensitive reabsorption of this sugar.